arXiv:math/9604209v1 [math.LO] 24 Apr 1996

Some interesting problems

Arnold W. Miller[1]

1. Analytic sets

1.1 (Mauldin) Is there a Σ^1_1 set X universal for Σ^1_1 sets which are not Borel? Suppose $B \in \Sigma^1_1$ and for every Borel A, $A \leq_W B$. Does this imply that for every Σ^1_1 A, $A \leq_W B$. (This refers to Wadge reducible.)

Answer: The first question was answered by Hjorth [41] who showed that it is independent.

1.2 A subset $A \subset \omega^\omega$ is compactly-Γ iff for every compact $K \subset \omega^\omega$ we have that $A \cap K$ is in Γ. Is it consistent relative to ZFC that compactly-Σ^1_1 implies Σ^1_1? (see Miller-Kunen [51], Becker [7])

1.3 ([51]) Does Δ^1_1 = compactly-Δ^1_1 imply Σ^1_1 = compactly-Σ^1_1?

1.4 (Prikry see [31]) Can $L \cap \omega^\omega$ be a nontrivial Σ^1_1 set? Can there be a nontrivial perfect set of constructible reals?

Answer: This is almost settled by Velickovic-Woodin [93].

1.5 (Ostaszewski [74]) Consider Telgarsky's game $G(T)$ where $T \subseteq 2^\omega$. Player I plays a countable cover of T Player II chooses one- say X_n. Player I wins iff $\cap\{cl(X_n) : n \in \omega\} \subseteq T$. It is known that

(a) Player I has winning strategy iff T is analytic.

(b) If there exists A an analytic subset of $cl(T)$ not Borel separated from T, then Player II has a winning strategy.

Is the converse of (b) true?

[1]April 1996. This is an update of my problem list [71]. I tried to include as many references as I could think of. If you know anything about these problems or could supply any missing references or corrections (missing attributions or misattributions), please let me know.

1.6 Does there exists an analytic set which is not Borel modulo Ramsey null? Same question for the ideal generated by closed measure zero sets.

1.7 (Sierpinski [85]) Does there exists an analytic set of reals E such that every (uncountable) analytic set of reals is the one-to-one continuous image of E?

1.8 (Jockusch 10-95) Let for $A \subseteq \omega$ let $D(A) = \{a - b : a, b \in A\}$. Is the set $\{D(A) : A \subseteq \omega\}$ Borel?

1.9 Suppose I is a σ-ideal generated by its Π^0_2 members. Then is it true that for any analytic set A either $A \in I$ or A contains a Π^0_3 set not in I? This is suggested by a theorem of Solecki [88] that says that for any σ-ideal I generated its closed members and analytic set A, either $A \in I$ or A contains a G_δ set not in I.

2. Axiom of Determinacy

2.1 Does AD imply that 2^{ω_1} is the ω_1 union of meager sets?

Answer: Yes, Becker [9].

2.2 Does AD imply that there does not exist ω_2 distinct Σ^1_2 sets?

2.3 Is there a hierarchy of Δ^1_2 sets?

2.4 Does AD imply every set is Ramsey?

Answer: Yes, if also assume $V = L[\mathbb{R}]$ for references see Kanamori [47] page 382.

2.5 (V. Delfino [12]) (Conjecture) If $f : 2^\omega \mapsto 2^\omega$ is Turing invariant ($x \equiv_T y \rightarrow f(x) \equiv_T f(y)$) then there exists z such that either for every $x \geq_T z$ $f(x) \geq_T x$ or there exist c such that for every $x \geq_T z$ $f(x) \equiv_T c$.

3. Combinatorial cardinals less than the continuum

3.1 (van Douwen [20]) If every ω_2 descending sequence in $P(\omega)$/finite has something beneath it is it true that every family of ω_2 sets with the IFIP has something beneath it? (does $\mathfrak{t} = \mathfrak{p}$).

3.2 (Hechler [36]) Let M be a countable transitive model of ZFC. Does there exists $M[f_n : n \in \omega]$ a generic extension with each $f_n \in \omega^\omega$ such that for all $n \in \omega$ f_n eventually dominates every element of

$$M[f_m : m > n] \cap \omega^\omega.$$

For something similar with Sacks forcing see Groszek [34].

3.3 (Dow) Does the following imply $p(\kappa)$: $\forall X, Y \subset [\omega]^\omega, |X|, |Y| < \kappa$ and $\forall A \in X, B \in Y$ $A \cap B$ finite, there exists $U \subset \omega$ such that for all $V \in (X \cup Y)$ $(U \cap V)$ is infinite iff $V \in X$

3.4 Can the least κ such that Indep(κ) fails have cofinality ω? Indep(κ) means that every family B of κ infinite subsets of ω there exists an infinite subset Z of ω such that for every $A \in B$, $|Z \cap A| = |Z \setminus A| = \omega$.

3.5 (Kunen [54]) Let $\mathfrak{m}$ be the smallest cardinal for which $\mathrm{MA}_{\mathfrak{m}}$ fails. Can we have $\omega_2 = cof(\mathfrak{m}) < \mathfrak{m}$?

3.6 (Scheepers 7-91) Is it consistent that $\aleph_\omega$ embeds into $(\omega^\omega, \leq^*)$ but not $\aleph_{\omega+1}$?

4. MAD families

4.1 (Roitman) Is it consistent that every maximal almost disjoint family in $[\omega]^\omega$ has cardinality greater than ω_1, but there exists a dominating family F in ω^ω of cardinality ω_1? For a related result see Shelah [80].

4.2 (van Douwen) CH implies there exist $F \subset \omega^\omega$ which is maximal with respect to eventually different functions which is also maximal with respect to infinite partial functions also. Is there always such a one? What is the cardinality of the smallest?

4.3 (Cook, Watson) Consider paths in $\omega \times \omega$. CH implies there is a MAD family of paths. Is there always one?

Answer: No, Steprans [86], still open for dimensions ≥ 3

4.4 (Milliken) A maximal almost disjoint family X is a separating family iff for all $Q \in [\omega]^\omega$

$$\{Q \cap P \mid P \in X\}$$

has size continuum or is finite. Are there always separating families?

4.5 (Erdös, Hechler [22]) Does MA plus the continuum is larger than $\aleph_{\omega+2}$ imply that there is no mad family on $\aleph_\omega$ of size $\aleph_{\omega+1}$?

4.6 (Kunen) Call $I \subset [\omega]^\omega$ an independent splitting family if I is independent (every finite boolean combination of elements of I is infinite) and splitting (for every $f : I \mapsto 2$ there does not exist an infinite X such that for every $A \in I$, $X \subset^* A^{f(A)}$, where $A^0 = A$ and $A^1 = \omega \setminus A$.) If CH or MA then there does exists an independent splitting family. In ZFC is there one?

4.7 (Fleissner) If there is a Luzin set, then is there a MAD family of size ω_1?

5. Forcing

5.1 (S. Friedman, R. David) Let $P_n = 2^{<\omega_n}$. Does forcing with $\Pi_{n\in\omega} P_n$ add a Cohen subset of $\omega_{\omega+1}$?

Answer: Yes, Shelah [83]

5.2 (Kunen) Force with perfect $P \subset 2^\omega$ such that for every $I \in [\omega]^\omega$ $\pi_I : P \mapsto 2^I$ does not have a countable range. Is ω_1 collapsed?

5.3 (van Douwen, Fleissner) Is it consistent with not CH that for P a c.c.c partial order of size continuum there exists a sequence G_α for $\alpha < \omega_1$ of P-filters such that for every dense set $D \subseteq P$ all but countably many of the G_α meet D.

Answer: No, Todorcevic [89]

5.4 Is there a Truss-like characterization of eventually different reals? How about infinitely equal reals? (see Truss [91] for characterization of dominating reals).

5.5 (Kunen [52]) Does there exists an ω_1 saturated σ-ideal in the Borel subset of 2^ω which is invariant under homeomorphisms induced by permutations of ω and different from the meager ideal, measure zero ideal, and the intersection ideal?

Answer: Partial answer Kechris-Solecki [48]

5.6 (van Mill) Is it consistent that every c.c.c. boolean algebra which can be embedded into $P(\omega)$/finite is σ-centered?

5.7 Suppose $M \subseteq M[f]$ are models of ZFC and for every $g \in \omega^\omega \cap M$ there exists infinitely many $n \in \omega$ such that $g(n) = f(n)$. Must there exists a real $x \in M[f]$ which is Cohen over M? (If there are two such infinitely equal reals, then there must be a Cohen real, see [66] and [3].)

6. Measure theory

6.1 (Mauldin, Grzegorek) Is it consistent that the continuum is RVM and all sets of reals of cardinality ω_2 have zero measure?

> Answer: No, apparently from the Gitik-Shelah Theorem (see Fremlin [30] 6F) it was deduced by Prikry and Solovay that if κ is real-valued measurable, then there are Sierpinski sets of all cardinalities less than κ.

6.2 (Fremlin) Can the cardinality of the least cover of the real line by measure zero sets have countable cofinality?

> Answer: Yes, Shelah [82]

6.3 (Erdös) For every sequence converging to zero does there exist a set of positive measure which does not contain a similar sequence? Falconer [25] has shown that if the sequence converges slowly enough there does exist such a set of positive measure. H.I. Miller [73] has shown the analogous statement for Baire category to be false. I showed that for every sequence there exist a partition of the reals into two sets neither of which contains a sequence similar to the given one.

6.4 (Erdös) Suppose for every $n \in \omega$ the set $A \cap [n, \infty)$ has positive measure. Must A contain arbitrarily long arithmetic progressions?

> Answer: Several mathematicians have pointed out this is trivial. Probably I misquoted Erdös. I scribbled it down after one of his talks when the universe was younger. To quote Just [44]: "The answer to 6.4 seems to be trivially 'yes', unless you want the differences to be integers; then the answer seems to be trivially 'no', unless you want the measure to be positive in EVERY interval, in which case the answer may not be so trivial. So, what should the problem really look like?"

6.5 Is it possible to have a Loeb-Sierpinski set of cardinality greater than ω_1? See Leth-Keisler-Kunen Miller [57] and Miller [69].

6.6 (Louveau) If a subset A of the plane has positive measure and contains the diagonal, then does there exist a set B in the line of positive outer measure such than B^2 is a subset of A?

Answer: According to Burke [11], Fremlin and Shelah proved this fails in the Cohen real model.

7. Borel hierarchies

7.1 Is it consistent that for every countable ordinal α there exists a Π^1_1 set of Baire order α? See Miller [62].

7.2 Is it consistent that for every uncountable separable metric space X there exists a X-projective set not Borel in X? See Miller [64],[70].

7.3 Is it consistent that the set of all Baire orders is the same as the set of even ordinals $\leq \omega_1$? See Miller [72].

7.4 Is it true that if X is a Q_α-set and Y is a Q_β-set and $2 \leq \alpha < \beta$ then $|X| < |Y|$? [62]

7.5 Does $\mathcal{R}^{\omega_2}_{\omega_1} = P(\omega_2 \times \omega_2)$ and $2^\omega = \omega_2$ imply that $2^{\omega_1} = \omega_2$? ($\mathcal{R}^{\omega_2}$ is the family of abstract rectangles in $\omega_2 \times \omega_2$ and the lower subscript is the level of the Borel hierarchy.)

7.6 Does $\mathcal{R}^{\omega_2}_{\omega} = P(\omega_2 \times \omega_2)$ imply that for some $n < \omega$ $\mathcal{R}^{\omega_2}_{n} = P(\omega_2 \times \omega_2)$?

7.7 Does $|X| = \omega_1$ imply that X is not a Q_ω-set?

7.8 (Mauldin) Is it consistent that there exists a separable metric space X of Baire order less than ω_1 (i.e. for some $\alpha < \omega_1$ every Borel subset of X is Σ^0_α in X) but not every relatively analytic set is relatively Borel?

7.9 Can the Borel hierarchy on cubes in $\mathcal{R}^3$ behave differently than the Borel hierarchy on rectangles in $\mathcal{R}^2$?

7.10 (Ulam [92]) Is there a separable metric space of each projective class order? (Σ^1_2-forcing?) See Miller [64],[70].

7.11 In the Cohen real model is there an uncountable separable metric space of Baire order 2? In the random real model are there any separable metric spaces of Baire order between 2 and ω_1?

Answer: Answered in Miller [72].

7.12 What can we say about hierarchy orders involving difference hierarchies or even abstract ω-boolean operations?

7.13 (Stone) Is it consistent to have a Borel map $f : X \mapsto Y$ where X and Y are metric spaces and f has the property that there is no bound less than ω_1 on

the Borel complexity of $f^{-1}(U)$ for $U \subseteq Y$ open? Fleissner [26] shows that it is consistent there is no such f using a supercompact.

7.14 (Ciesielski-Galvin [15]) Let $P_2(\kappa)$ be the family of all cylinder sets in κ^3 (where cylinder means $A \times B$ where $A \subseteq \kappa$ and $B \subseteq \kappa^2$ or anything that could be obtained like this by permuting the three coordinates.) Is it consistent that the σ-algebra generated by $P_2(\mathfrak{c}^{++})$ is equal to all subsets of $(\mathfrak{c}^{++})^3$?

7.15 (Ciesielski) Suppose every subset of $\omega_2 \times \omega_2$ is in the σ-algebra generated by the abstract rectangles. Does this continue to hold after adding ω_1-Cohen reals?

7.16 (Fleissner [27]) If X is a Q-set of size ω_1 then is X^2 a Q-set? (Not necessarily true if for X of cardinality ω_2.)

8. Involving ω_1

8.1 (Jech-Prikry [42]) Is it consistent that there exists a family $F \subset \omega^{\omega_1}$ of cardinality less than 2^{ω_1}, such that for every $g \in \omega^{\omega_1}$ there exist $f \in F$ such that for every $\alpha < \omega_1, g(\alpha) < f(\alpha)$.

8.2 (Frankiewicz [28]) Is it consistent that $\beta\omega \setminus \omega$ is homeomorphic to $\beta\omega_1 \setminus \omega_1$?

8.3 Is it consistent to have CH, $2^{\omega_1} > \omega_2$, and there exists $F \subset [\omega_1]^{\omega_1}$ of cardinality ω_2 such that for every $A \subset \omega_1$ there exists $B \in F$ such that $B \subseteq A$ or $B \cap A = \emptyset$?

8.4 (Kunen) Is it consistent to have $2^{\omega_1} > \omega_2$ and there exists $F \subset [\omega_1]^{\omega_1}$ of cardinality ω_2 such that for every uncountable $A \subset \omega_1$ there exists $B \in F$ such that $B \subseteq A$?

8.5 (Kunen) Is it consistent to have a uniform ultrafilter on ω_1 which is generated by fewer than 2^{ω_1} sets?

8.6 (Prikry [78]) Is it consistent there exists an ω_1 generated ideal J such that $P(\omega_1) = P(\omega)/J$?

8.7 (Comer) If C and D are homeomorphic to 2^{ω_1} then is $C \cup D$? (Say if both are subsets of 2^{ω_1}.)

8.8 (Nyikos) If $C \times D$ is homeomorphic to 2^{ω_1} then must either C or D be homeomorphic to 2^{ω_1}?

8.9 (CH) Let n(X) be the cardinality of the smallest family of meager sets which cover X. Can the cof(n($(2^{\omega_1})_\delta$)) (G_δ-topology) be ω or ω_1?

8.10 (Velickovic) Is it consistent that every Aronszajn line L contains a Countryman type?

8.11 Does PFA imply that any two Aronszajn types contain uncountable isomorphic subtypes?

9. Set theoretic topology

9.1 Is it consistent to have no P-points or Q-points? A P-point is an ultrafilter U on ω with the property that every function $f : \omega \mapsto \omega$ is either constant or finite-to-one on an element of U. A Q-point is an ultrafilter U on ω with the property that every finite-to-one function $f : \omega \mapsto \omega$ is one-to-one on an element of U. Shelah [79] showed it is consistent there are no P-points and Miller [63] showed that it is consistent there are no Q-points. Roitman and Taylor showed that if the continuum is $\leq \omega_2$, then there must be a P-point or a Q-point.

9.2 (M.E. Rudin) Is there always a small Dowker space?

Answer: Yes?, Balogh [2], Kojman-Shelah [49]

9.3 (Charlie Mills) In infinite dimensional Hilbert space is a sphere coverable by fewer than continuum other spheres?

9.4 Is it consistent that ω^{ω_1} is pseudonormal? (Pseudonormal means disjoint closed sets can be separated if at least one is countable.)

9.5 (van Douwen) Is it consistent to have $c(U(\omega_1)) < d(U(\omega_1)$? (c is celluarity, d is density and U is uniform ultrafilters.)

9.6 Is the box product of countably many copies of the unit interval coverable by countably many zero dimensional sets?

9.7 (Hansell) Is there a non-zero-dimensional Q-space? Can there be a nonzero dimensional metric space in which every subset is G_δ?

9.8 (Bing) Suppose D_n a subset of the plane is homeomorphic to a disk and for every $n \in \omega$ $D_{n+1} \subseteq D_n$, then does $\cap_{n\in\omega} D_n$ have the fixed point property?

9.9 (Sikorski see [55]) Does there exist two closed 0-dimensional nonhomeomorphic subsets of the plane such that each is homeomorphic to an open subset of the other?

9.10 (Ancel 11-93) Is there a separable Hausdorff space in which every basis has cardinality 2^{2^c}? (c = the cardinality of the set of real numbers.)

9.11 (Gulko 1995) Is there a model of ZFC in which there is a maximal almost family M on ω such that for any point $x \in \beta\omega \setminus \omega$ there exists a countable subfamily of M such that x is in its closure.

10. Model Theory

10.1 (Vaught) Does every countable first order theory have countably many or continuum many countable models up to isomorphism? How about for universal theories of a partial order? See Becker [8].

10.2 (Martin) Show that if T is a countable first order theory with fewer than continuum many countable models up to isomorphism, then every countable model of T has an isomorphism class which is at most $\Sigma^0_{\omega+\omega+1}$.

10.3 (Caicedo [13]) Does every theory in $L_{\omega_1,\omega}$ have an independent axiomatization?

10.4 Does V=L imply there exists a complete theory T such that

$$\{\alpha : L_\alpha \models T\}$$

is an unbounded subset of ω_1?

Answer: Yes, Hjorth [37].

10.5 (Miller [67]) Are there any properly $\Sigma^0_{\lambda+1}$ isomorphism classes for λ a countable limit ordinal?

Answer: Yes, Hjorth [38]

10.6 Is there a theory with exactly ω_1 rigid countable models up to isomorphism? (same for minimal models)

Answer: Yes, for minimal models, Hjorth [39]. Yes, for rigid models, Hjorth [40]

10.7 (Baldwin) Are there continuum many complete ω-stable theories in a finite language? (I don't know if this is still open?)

10.8 (Miller-Manevitz [60]) Is it consistent that there exists a model of ZFC, M, such the unit interval of M is Lindelof and ω^M is ω_1-like?

10.9 (Miller [67]) Does there exists a pseudo-elementary class in the language of one unary operation with exactly ω_1 nonisomorphic countable models? (There is pseudo $L_{\omega_1,\omega}$ class.)

10.10 (Mati Rubin) Does there exists an embedding of the rationals into themselves such that no between function is elementarily extendable?

10.11 Duplicate of 10.6

10.12 (suggested by Fuhrken [32] [24]) Can we have a model with exactly one undefinable $L_{\omega_1,\omega}$ element?

10.13 Can we have a complete first order theory T with models of size $\aleph_{2n}$ for $n < \omega$ (but not of size $\aleph_{2n+1}$) and $\aleph_\omega < c$?

11. Special subsets of the real line

11.1 (Mauldin, Grzegorek) Is it consistent that every universally measurable set has the Baire property? See Corazza [17].

11.2 (Mauldin) Are there always $> \mathfrak{c}$ universally measurable sets? (same for restricted Baire property)

11.3 (Galvin) Does every Sierpinski set have strong first category?

> Answer: Bartoszynski-Judah [4] showed that it is consistently yes. Yes, Pawlikowski [75].

11.4 (Galvin, Carlson) Is the union of two strong first category sets a set of strong first category?

> Answer: Not necessarily, Bartoszynski-Shelah [5]

11.5 Does there exists a perfectly meager $X \subseteq \mathbb{R}^n$ which is not zero-dimensional?

11.6 (Kunen) Is it consistent that for every uncountable $X \subseteq \mathbb{R}$ there exists a measure zero set M such that $X + M$ has positive outer measure? See Erdos-Kunen-Mauldin [23].

11.7 (Sierpiński [84]) A set of reals X is a J-set iff for uncountable $Y \subseteq X$ there exist a perfect $P \subseteq X$ such that $P \cap Y$ is uncountable. If we assume CH, then a set is a J-set iff it is σ-compact. Is it consistent with not CH that J-set = σ-compact?

11.8 (Fremlin-Miller [29]) Is there always an uncountable subset of the reals which is hereditary with respect to property M?

> Answer: Yes, see [45]

11.9 Consider the three non c.c.c ideals: $(s)_0$-sets, Ramsey null sets, and σ-compact sets. What can one say about the properties of add, cov, non, and cof? (add = additivity of ideal, cov = smallest cardinality of a cover

of reals by subset of ideal, non = smallest cardinality of a set of reals not in ideal, cof = cofinality = smallest cardinality of a family of sets in ideal which has the property that every set in ideal is covered by some element of the family.)

11.10 Consider the notion of Laver null sets. This is defined analogously to Ramsey null sets, but use Laver forcing instead of Mathias forcing. The analogue of Galvin-Prikry Theorem is true here. What other results also go thru? What ideals arise from other notions of forcing? What about Silver forcing? What notions of forcing arise from infinite combinatorial theorems? (For example, Carlson's infinite version of the Hales-Jewett theorem [14].)

11.11 (Judah-Shelah [43]) Is the Borel conjecture plus the existence of a Q-set consistent?

11.12 (Daniel, Gruenhage). Given a set of reals X and ordinal α let $G_\alpha(X)$ be the game of length α played by two players: point picker and open. At each play of the game point picker picks a real and open responds with an open set including the real. Point picker wins a run of the game if at the end the open sets chosen cover X. The order of X is the least ordinal for which point picker has a winning strategy. What orders are possible? Daniel and Gruenhage have examples of order ωn assuming CH.

11.13 (Komjath, see[87]) Suppose every set of reals of size ω_1 has measure zero. Then does every ω_1 union of lines have planar measure zero? (Dually) Suppose the real line is union of ω_1 measure zero sets. Then does there exists ω_1 measure zero subsets of the plane such that every line is contained in one of them?

11.14 (Zhou 3-93) Does every set of size ω_1 is a Q-set imply that $\mathfrak{p} > \omega_1$. For γ-sets it is true.

12. Quasiorder theory

12.1 Is there a Borel version of Fraisse's conjecture? Are the Borel linear orderings well-quasiordered under embedding?

Answer: Yes, Louveau and St-Raymond [59] assuming large parts of AD.

12.2 (Laver) Is it consistent that the set of Aronszajn trees is well-quasi-ordered under embeddability? See Laver [56] and Corominas [18].

12.3 (Kunen) Is the set of all better-quasi-ordered binary relations on ω a proper Π^1_2 set?

Answer: Yes, Marcone [61]

12.4 Suppose $(Q, \leq)$ is a recursive quasi-order. Is it true that Q is BQO iff $Q^{<\omega_1^{ck}}$ is WQO?

12.5 (Kunen) Suppose $(Q, \leq)$ is a recursive well quasi-order. Does $Q^\omega/\equiv$ have a recursive presentation?

12.6 Suppose every set is Ramsey and $f : [\omega]^\omega \mapsto$ ORD. Then does there exist $X \in [\omega]^\omega$ such that the image of $[X]^\omega$ under f is countable? See Louveau-Simpson [58] and Aniszcyk-Frankiewicz-Plewik [1].

12.7 Is finite graphs under homeomorphic embedding WQO?

12.8 Is the witness lemma true for LIN(Q) or TREE(Q)?

12.9 Is there an ω_1-descending sequence of countable posets (under embedding) each of which is the union of two chains? (Kunen, Miller: There is an ω_1-descending sequence of countable posets. Kunen: There is an infinite antichain of finite posets each of which is the union of two chains.)

12.10 Is there a parameterized version of Carlson's theorem? See Carlson [14] and Pawlikowski [76].

13. not AC

13.1 (M. Bell) Does ZF imply that for every family of nonempty sets there exists a function assigning to each set in the family a compact Hausdorff topology? (Motivation: AC is equivalent to this principal plus every Tychonov product of compact Hausdorff spaces is compact.)

Answer: No, Todorcevic [90] Just [44]

13.2 (Dow 88) Does Stone's theorem on metric spaces (every metric space is paracompact) require AC? It is known that ZF implies that ω_1 with the order topology is not metrizable.

Answer: Yes, Watson see [33].

14. Recursion theory

14.1 Does there exist a non-trivial automorphism of the Turing degrees? (Re degrees?)

Answer: Yes for both? announced by Cooper [16]

14.2 (jockusch) Does there exists a DNR of minimal Turing degree? (DNR means diagonally non recursive: $f \in \omega^\omega$ and for all $e \in \omega$, $f(e) \neq \{e\}(e)$.)

15. Miscelleneous

15.1 (Sierpiński) Is there a Borel subset of the plane which meets every line in exactly two points? (Mauldin) Must such a set be zero dimensional?

> Answer: Davies has shown such a set cannot be Σ^0_2 and Mauldin has shown such a set must be disconnected. Miller [68] showed that if V=L then there does exist a Π^1_1 subset of the plane which meets every line in exactly two points. Kulesza [50] showed that any two point set must be zero dimensional.

15.2 (Cichon) Is it consistent to have that the real line is the disjoint union of ω_2 meager sets such that every meager set is contained in a countable union of them?

> Answer: No, Brendle [10]

15.3 (Juhasz) Does club imply there exist a Souslin line?

> Answer: No, Dzamonja-Shelah [21]

15.4 (Ulam [92]) Does there exist a set D dense in the plane such that the distance between any two points of D is rational?

15.5 (Miller [65]) Suppose the continuum is greater than ω_2, then does there exists a set of reals of cardinality the continuum which cannot be mapped continuously onto the unit interval?

15.6 Is it consistent that there exists $x \in 2^\omega$ such that $V = L[x] \neq L$ and a continuous onto function $f : L \cap 2^\omega \mapsto V \cap 2^\omega$?

15.7 (Price [77]) Is it consistent there is no Cech function?

15.8 (Kunen) Does the consistency of an elementary embedding of M into V imply the consistency of a measurable cardinal?

15.9 (Erdös) Without CH can you partition the plane into countably many pieces so that no piece contains an isoceles triangle? See Kunen [53].

15.10 Is there a Borel version of Hall's marriage theorem? As for example, the Borel-Dilworth Theorem [35].

15.11 (Davies [19]) Assuming CH for every $f : \mathbb{R}^2 \mapsto \mathbb{R}$ there exists $g_n, h_n : \mathbb{R} \mapsto \mathbb{R}$ such that

$$f(x, y) = \Sigma_{n\in\omega} g_n(x)h_n(y)$$

Does this imply CH?

15.12 (Mauldin) CH implies that for every $n \geq 3$ there exists a 1-1 onto function $f : \mathbb{R}^n \mapsto \mathbb{R}^n$ which maps each circle onto a curve which is the union of countably many line segments. Is CH necessary?

15.13 (Kunen) Can there be a Souslin tree $T \subseteq 2^\kappa$ such that for all $\alpha < \kappa$ the T_α contains all except at most one of the α branches thru $T_{<\alpha}$. Here κ is the first Mahlo or weakly Mahlo.

15.14 (Baumgartner [6]) Is it consistent that any two ω_2 dense sets of reals are order isomorphic?

15.15 (S. Kalikow [46]) For any set X define for $x, y \in X^\omega$, $\quad x =^* y$ iff for all but finitely many $n \in \omega$, $\quad x(n) = y(n)$. X has the discrete topology and X^ω the product topology. Is it consistent that there exists a map $f : \omega_2^\omega \mapsto 2^\omega$ which is continuous and for every $x, y \in {\omega_2}^\omega$, $\quad x =^* y$ iff $f(x) =^* f(y)$. (Kalikow: yes for ω_1 in place of ω_2.)

Answer: Yes, Shelah [81].

15.16 (unknown 1-92) According to Erdos, Sylvestor proved that given finitely many points F in the plane not all collinear, there exists a line L which meets F in exactly two points. $F = \mathbb{Z} \times \mathbb{Z}$ is an obvious infinite counterexample. Does there exists a counterexample which is a convergent sequence? countable compact set?

References

[1] B.Aniszcyk, R.Frankiewicz, and S.Plewik, Remarks on (s) and Ramsey-measurable function, Bulletin de L'Academie Polonaise des sciences, 35(1987), 479-485.

[2] Z.Balogh, A small dowker space in ZFC, preprint 6-95. where???

[3] T.Bartoszynski, Combinatorial aspects of measure and category, Fundamenta Mathematicae, 127(1987), 225-239.

[4] T.Bartoszynski, H.Judah, On Sierpinski sets, Proceedings of the American Mathmatical Society, 108(1990), 507-512.

[5] T.Bartoszynski, S.Shelah, consistent to have union of strong first category not strong first category, announced 95. where???

[6] J.Baumgartner, All $\aleph_1$-dense sets of reals can be isomorphic, Fundamenta Mathematicae, 79(1973), 101-106.

[7] H.Becker, Analytic sets from the point of view of compact sets, Mathematical Proceedings of the Cambridge Philosophical Society, 99(1986), 1-4.

[8] H.Becker, The topological Vaught's conjecture and minimal counterexamples, Journal of Symbolic Logic, 59(1994), 757-784.

[9] H.Becker, Solution to a particularly interesting problem of Arnie Miller, handwritten 1-94. where???

[10] J.Brendle, Nicely generated and chaotic ideals, preprint < 95. where???

[11] M.Burke, A theorem of Friedman on rectangle inclusion and its consequences, note dated March 7,1991.

[12] Cabal Seminar 76-77,77-79,79-81,81-85, Lecture Notes in Mathematics, 689,839,1019,1333, Springer-Verlag.

[13] X.Caicedo, Independent sets of axioms in $L_{\kappa\alpha}$, Canadian Mathematical Bulletin, 24(1981), 219-223.

[14] T.Carlson, Some unifying principles in Ramsey Theory, Disc. Math. 68 (1988) 117-169.

[15] K.Cielsielski, F.Galvin, Cylinder problem, Fundamenta Mathematicae, 127(1987), 171-176.

[16] B.Cooper, Automorphism of the degrees, announced 95. where???

[17] P.Corazza, Ramsey sets, the Ramsey ideal, and other classes over $\mathbb{R}$, Journal of Symbolic Logic, 57(1992), 1441-1468.

[18] E.Corominas, On better quasiordering countable trees, Discrete Mathmatics, 53(1985), 35-53.

[19] R.O.Davies, Representation of functions of two variables as sums of rectangular functions I, Fundamenta Mathematicae, 85(1974), 177-183.

[20] E.van Douwen, The integers and topology, in **Handbook of set theoretic topology**, ed by K.Kunen and J.Vaughan, North-Holland, (1984), 111-167.

[21] M.Dzamonja, S.Shelah, $\clubsuit$ does not imply the existence of a Suslin tree, preprint 4-96, DjSh 604. where???

[22] P.Erdos, S.Hechler, On maximal almost-disjoint families over singular cardinals, Colloquia Mathematica Societatis Janos Bolyai, 10(1973), Infinite and finite sets, Keszthely Hungary, 597-604.

[23] P.Erdos, K.Kunen, and R.Mauldin, Some additive properties of sets of real numbers, Fundamenta Mathematicae, 113 (1981),187-199.

[24] A.Ehrenfeucht, G.Fuhrken, On models iwth undefinable elements, Math. Scand. 28(1971), 325-328.

[25] K.Falconer, On a problem of Erdos on sequences and measurable sets, Proceedings of the American Mathematical Society, 90(1984), 77-78.

[26] W.Fleissner, An axiom for nonseparable Borel theory, Transactions of the American Mathematical Society, 251(1979), 309-328.

[27] W.Fleissner, Squares of Q sets, Fundamenta Mathematicae, 118(1983), 223-231.

[28] R.Frankiewicz, To distinguish topologically the spaces m^*, Bulletin Academie Polonaise Science, 25(1977), 891-893.

[29] D.Fremlin and A.Miller, On some properties of Hurewicz, Menger, and Rothberger, Fundamenta Mathematicae, 129(1988), 17-33.

[30] D.Fremlin, Real-valued-measurable cardinals, in **Israel Mathematical Conference Proceedings**, ed by H.Judah, 6(1993), 151-304.

[31] H.Friedman, One hundred and two problems in mathematical logic, Journal of Symbolic Logic, 40(1975), 113-129.

[32] G.Fuhrken, A model with exactly one undefinable element, Colloquium Mathmaticum, 19(1968), 183-185.

[33] C.Good, I.J.Tree, Continuing horrors of topology without choice, Topology and its applications 63(1995), 79-90.

[34] M.Groszek, ω_1^* as an initial segment of c-degrees, Journal of Symbolic Logic, 59(1994), 956-976.

[35] L.Harrington, D.Marker, S.Shelah, Borel orderings, Transactions of the American Mathematical Society, 310(1988), 293-302.

[36] S.Hechler, On the existence of certain cofinal subsets of ${}^{\omega}\omega$, in **Axiomatic Set Theory**, proceedings of symposia in pure mathematics, vol 13 part 2, ed by T.Jech, (1974), 155-174.

[37] G.Hjorth, email message 11-94. where???

[38] G.Hjorth, An orbit that is exactly $\Sigma^0_{\lambda+1}$, handwritten note, 2-95. where???

[39] G.Hjorth, On $\aleph_1$ many minimal models, handwritten note, 12-94. where???

[40] G.Hjorth, On $\aleph_1$ many rigid models, handwritten note, 4-96. where???

[41] G.Hjorth, Universal co-analytic sets, preprint 12-94. where???

[42] T.Jech, K.Prikry, Ideals over uncountable sets: applications of almost disjoint functions and generic ultrapowers, Memoirs of the American Mathematical Society, 18 no 214, (1979).

[43] H.Judah, S.Shelah, Q-sets do not necessarily have strong measure zero, Proceedings of the American Mathematical Society, 102(1988), 681-683.

[44] W.Just, email 12-93. where???

[45] W.Just, A.Miller, M.Scheepers, and P.Szeptycki, The combinatorics of open covers (II), preprint 1995. where???

[46] S.Kalikow, Sequences of reals to sequences of zeros and ones, Proceedings of the American Mathematical Society, 108(1990), 833-837.

[47] A.Kanamori, **The higher infinite**, Springer-Verlag, (1994).

[48] A.Kechris, S.Solecki, Approximation of analytic by Borel sets and definable countable chain conditions, preprint 95. where???

[49] M.Kojman, S.Shelah, A ZFC Dowker space in $\aleph_{\omega+1}$: an application of pcf theory to topology, preprint 1996. where???

[50] J.Kulesza, A two point set must be zero dimensional, Proceedings of the American Mathematical Society, 116(1992), 551-553.

[51] K.Kunen and A.Miller, Borel and projective sets from the point of view of compact sets, Mathematical Proceedings of the Cambridge Philosophical Society, 94(1983), 399-409.

[52] K.Kunen, Random and Cohen reals, in Handbook of Set Theoretic Topology, North-Holland, (1984), 887-911.

[53] K.Kunen, Partitioning Euclidean space, Mathematical Proceedings of the Cambridge Philosophical Society, 102(1987), 379-383.

[54] K.Kunen, Where MA first fails, Journal of Symbolic Logic, 53(1988), 429-433.

[55] K.Kuratowski, On a topological problem connected with the Cantor-Berstein theorem, Fundamenta Mathematicae, 37(1950), 213-216

[56] R.Laver, Better-quasi-orderings and a class of trees, in Studies in Foundations and Combinatorics, Advances in Mathematics Supplementary Studies (1978) 31-48.

[57] S.Leth, J.Keisler, K.Kunen and A.Miller, Descriptive set theory on a hyperfinite set, Journal of Symbolic Logic, 54(1989), 1167-1180.

[58] A.Louveau and S.Simpson, A separable image theorem for ramsey mappings, Bulletin de L'Academie Polonaise des sciences, 30(1982), 105-108.

[59] A.Louveau and J. Saint Raymond, On the quasi-ordering of Borel linear orders under embeddability, Journal of Symbolic Logic 55(1990), 537-560.

[60] L.Manevitz and A.Miller, Lindelöf models of the reals: solution to a problem of Sikorski, Israel Journal of Mathematics, 45(1983), 209-218.

[61] A.Marcone, The set of better-quasi-orderings is Π^1_2, Mathematical Logic Quarterly, 41(1995), 373-383.

[62] A.Miller, On the length of Borel hierarchies, Annals of Math Logic, 16(1979), 233-267.

[63] A.Miller, There are no Q-points in Laver's model for the Borel conjecture, Proceedings of the American Mathematical Society, 78(1980), 103-106.

[64] A.Miller, Generic Souslin sets, Pacific Journal of Mathematics, 97(1981), 171-181.

[65] A.Miller, Mapping a set of reals onto the reals, Journal of Symbolic Logic, 48(1983), 575-584.

[66] A.Miller, A characterization of the least cardinal for which the Baire category theorem fails, Proceedings of the American Mathematical Society, 86(1982), 498-502.

[67] A.Miller, The Borel classification of the isomorphism class of a countable model, Notre Dame Journal of Formal Logic, 24(1983), 22-34.

[68] A.Miller, Infinite combinatorics and definability, Annals of Pure and Applied Mathematical Logic, 41(1989), 179-203.

[69] A.Miller, Set theoretic properties of Loeb measure, Journal of Symbolic Logic, 55(1990), 1022-1036.

[70] A.Miller, Projective subsets of separable metric spaces, Annals of Pure and Applied Logic, 50(1990), 53-69.

[71] A.Miller, Some interesting problems, Set Theory of the Reals, ed Haim Judah, Israel Mathematical Conference Proceedings, vol 6 (1993), 645-654, American Math Society.

[72] A.Miller, **Descriptive Set Theory and Forcing: how to prove theorems about Borel sets the hard way**, Lecture Notes in Logic 4(1995), Springer-Verlag.

[73] H.I.Miller, Some results connected with a problem of Erdos II, Proceedings of the American Mathematical Society, 75(1979), 265-268.

[74] Ostaszewski, email 9-92. where???

[75] J.Pawlikowski, Every Sierpinski set is strongly meager, preprint < 95. where???

[76] J.Pawlikowski, Parametrized Ellentuck Theorem, Topology and its Applications, 37(1990), 65-73.

[77] R.Price, On a problem of Čech, Topology and Its Applications, 14(1982), 319-329.

[78] K.Prikry and A.Miller, When the continuum has cofinality ω_1, Pacific Journal of Mathematics, 115(1984), 399-407.

[79] S.Shelah, **Proper Forcing**, Lecture Notes in Math, 940 (1982), Springer-Verlag.

[80] S.Shelah, On cardinal invariants of the continuum, Collection: Axiomatic set theory Boulder, Colo., 31(1983), 183-208 Contemp. Math., AMS.

[81] S.Shelah, On a problem of Steve Kalikow, preprint (1995) Sh590. where???

[82] S.Shelah, Covering of the null ideal may have countable cofinality, preprint 6-95, Sh592. where???

[83] S.Shelah, Embedding Cohen algebras using pcf theory, preprint 7-95, Sh595. where???

[84] W.Sierpinski, **Hypothese du Continu**, Monografje Matematyczme, Warszawa-Lwow, 1934.

[85] W.Sierpinski, Problem 70, Fundamenta Mathematicae, 26(1936), 334.

[86] J.Steprans, Almost disjoint families of paths in lattice grids, Topology Proceedings 16(1991) 185-200.

[87] J.Steprans, Cardinal invariants associated with Hausdorff capacities, preprint 6-94. where???

[88] S.Solecki, Covering analytic sets by families of closed sets, Journal of Symbolic Logic, 59(1994), 1022-1031.

[89] S.Todorcevic, Remarks on Martin's axiom and the continuum hypothesis, Canadian Journal of Mathematics, 43 (1991), 832-851.

[90] S.Todorcevic, handwritten note 8-91. where???

[91] J.Truss, Sets having calibre $\aleph_1$, Logic Colloquium 76, North-Holland, Amsterdam, 1977, 595-612.

[92] S.Ulam, **Problems in Modern Mathematics**, Wiley, New York, 1964.

[93] B.Velikovic, H.Woodin, Complexity of reals of inner models of set theory, preprint 95. where???

Arnold W. Miller
University of Wisconsin
Department of Mathematics, Van Vleck Hall
480 Lincoln Drive
Madison, WI 53706
miller@math.wisc.edu
http://www.math.wisc.edu/∼miller